


Vaccination and polio eradication

Prof Nicholas Grassly & Dr Tara Mangal



Poliomyelitis: the virus

- ssRNA enterovirus
- Three distinct genetic types with no (very limited) cross immunity (1,2,3)
- Humans are only known natural host



Poliomyelitis: an ancient disease




Egypt ~ 1500 BC

Prodromal non-paralytic phase leads to acute flaccid paralysis (AFP) in about 1 in 200 cases


In approximately 1 in 10 AFP cases the brain stem is affected, leading to paralysis of breathing muscles and death

Post-polio syndrome (PPS) - polio patients who recovered well due to re-innervation may experience muscular problems in later life

Pathogenesis




Natural history of infection (1)



- Incubation period average 16.5 days

- Around 3-5 days after exposure virus can be detected in faeces, blood and throat



- Virus is detected in the faeces for on average 4-5 weeks *after onset of symptoms*, but for only c. 2 weeks in the throat

Natural history of infection (2)


Risk of AFP increased for:

Age c. 5-15 yrs

Women at older ages,
men at younger ages

Bigger viral dose

Subtype (1>3>2)



Following infection long-lasting homologous humoral immunity occurs (e.g. eskimos)

Mucosal immunity may wane, facilitating transmission without disease

Limited evidence for short-lived heterotypic immunity

Modes of transmission

1. Droplet
2. Fomite and direct contact
3. Faecal oral

Relative importance of transmission route depends on context and may have implications for dynamics e.g. faecal-oral probably less important in developed world

Descriptive epidemiology

- Pre-vaccination
 - Developing countries
 - Endemic disease
 - Most (>90%) children infected by 4 years of age
 - Industrialized countries
 - Epidemic polio with increasingly severe epidemics perhaps partly due to older average age at infection (e.g. early 1950s average age at infection in US 5-9 years, cf. <5 years in the 1916 epidemic) but also increased diagnosis and care (Anderson and May 1991)
- Post-vaccination
 - Developing countries
 - Ongoing polio eradication efforts in Africa and South East Asia
 - Industrialized countries
 - Disease eradicated

Increasing number of epidemics identified

- Increasingly frequent outbreaks in early 20th century
- Franklin D Roosevelt infected in 1921
- Founded the March of Dimes in 1938 (originally called National Foundation for Infantile Paralysis)
- “Great Race” for a vaccine began




Inactivated Polio Vaccine (IPV)

- First developed by Salk in 1952 by prolonged treatment of virus with formalin
- Injected
- Associated with ‘Cutter’ incident
- Current preparation is through purification and concentration of inactivated wild poliovirus
- Induces good ‘humoral’ immunity against paralysis
- Much more limited impact on mucosal immunity against infection

Oral Polio Vaccine (OPV) (1)

- Developed by Sabin in 1962
- Attenuated by passage through different host cells
- Stability of attenuated form assessed empirically by inoculating monkeys
- However, only a few mutations separate OPV strains from wildtype (particularly types 2 and 3)
- Vaccine Associated Paralytic Poliomyelitis (VAPP) occurs
 - 1 in 750,000 receiving 1st dose
 - 1 in 12 million receiving 2nd dose




Oral Polio Vaccine (OPV) (2)

- Trivalent and monovalent live virus vaccines have been licensed
- Vaccine is infectious and may secondarily immunise household contacts of vaccinee (e.g. un-immunized gypsy populations in Europe have high levels of serum neutralising antibodies to polio types 1,2 and 3 probably due to infection with OPV)
- With trivalent vaccine >95% seroconvert to each poliovirus type after 3 doses in developed countries
- Typically only ~70-80% (or less) seroconvert to type 1 & 3 after 3 or 4 doses in developing countries due to interference by other infections and diarrhoeal disease
- Seroconversion is also associated with induction of good mucosal immunity against infection

Which vaccine to use?

- OPV cheaper, easier to administer, secondarily immunises contacts of vaccinees, protects against infection (mucosal immunity)
- Majority of countries switched to OPV from IPV during the 1960s
- Wealthy countries now switching back to IPV




Eradication - Justification

- No natural non-human reservoir
- Effective vaccine - lifelong serum immunity
- Although vertical programme, can have positive impact on health services
 - e.g. eradication of smallpox (1958-1980) was basis of EPI
 - (failed) eradication initiative for yellow fever (1915-1977) led to first national administrative health systems in many countries
- Governments of the world committed in 1988 with initial aim to eradicate disease by 2000, revised to 2005
- Would be second infectious disease to be eradicated after smallpox




Vaccination and herd immunity

- Direct (individual) effect
 - vaccinee protected against infection
- Indirect (population) effect
 - reduced force of infection leading to ‘herd immunity’
- Effective reproductive number R



$$p_c = \frac{R_0 - 1}{R_0} = 1 - \frac{1}{R_0}$$

Critical vaccination threshold



$$p_c = 1 - \frac{1}{R_0}$$

Eradication - challenges


- More difficult than Smallpox eradication since
 - Need more than 1 dose of vaccine
 - (3 in developed countries with good sanitary conditions, 8 or more in countries with poor sanitation)
 - AFP may be caused by other enteroviruses or infections; therefore symptoms less specific
 - Larger R_0 ? (variable estimates e.g. see Anderson and May 1991)
 - Bigger more mobile population (7 vs 4 billion)
- Also
 - Expensive (US\$6 billion external expenditure to date)
 - Diversion of resources?
 - Achievable?

Global polio surveillance


- Monitoring of AFP cases
- Need to distinguish polio derived AFP from other causes (Guillain-Barré syndrome, trauma, transverse myelitis and other enterovirus infections)
- >30,000 AFP cases now undergo clinical, epidemiologic and virologic investigation every year by a network of laboratories co-ordinated by CDC and WHO
- Any wildtype polio viruses are sequenced to allow identification of origin of outbreak
- But 'silent spread' can occur, since only 1 in 200 infected develop AFP

Surveillance indicators, polio endemic regions

Non-polio AFP rate



Adequate stool collection rate



March - August 2012

The boundaries and names shown and the designations used on this map do not imply the expression of any opinion whatsoever on the part of the World Health Organization concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. Dotted lines on maps represent approximate border lines for which there may not yet be full agreement.

© WHO 2012. All rights reserved

Data in WHO/HQ as of 02 Oct 2012

Massive campaigns




NIDs Launch
Nigeria

- Cuba first country to have a NID in 1963
- e.g. in October 2004 during four day synchronised NIDs 100 million OPV doses given across half of Africa
- Type 2 has been eradicated globally (last detected in Aligarh, India 1999)


Global polio eradication: end stages

Reduction in incident cases of paralytic polio from 1000 per day in 1988 to 5 per day in 2006

Stopped poliovirus transmission in the Americas, Western Pacific and Europe





Global situation 2008



- Wild virus type 1
- Wild virus type 3
- Endemic countries
- ▨ Case or outbreak following importation in last 6 months

Polio persistence in India



Poliovirus	Location	Cases	Matches	Vaccine efficacy (95% CI)
Type 1	rest of India	1512	361	21 (15 - 27)
	Bihar	387	158	18 (9 - 26)
	Uttar Pradesh	2522	1108	9 (6 - 13) **
Type 3	rest of India	221	79	21 (8 - 33)
	Bihar	136	53	22 (4 - 36)
	Uttar Pradesh	847	342	9 (3 - 15)


** significantly different from rest of India, $p < 0.01$

New strategies


- Type 2 wild poliovirus eradicated in 1999 and type 3 transmission geographically limited
- New high potency (10^6 CCID₅₀/dose) monovalent type 1 vaccine developed and licensed through extraordinary public-private partnership in less than a year 2004-5
- Potentially more efficacious since no interference between vaccine strains
- First used in India and Egypt in 2005
- Estimated field efficacy in India and Nigeria 3-4 times that of the trivalent vaccine against paralytic type 1 poliovirus



Bivalent OPV



Non-inferiority and superiority assessments after a cumulative two-dose schedule of vaccines (Sutter et al. *Lancet* 2010)



Estimated proportion of children with non-polio AFP under two years old in Nigeria protected by vaccination against poliomyelitis caused by serotype 1 (red lines) or 3 (green lines) poliovirus. These age-standardised estimates exclude the effect of natural exposure to wild-type or vaccine poliovirus and protection offered by maternal antibodies.

Vaccine efficacy estimates are derived from analysis of 2001-2011 data from Nigeria.

	Serotype 1	Serotype 3
tOPV	18 (15-22)	19 (12-24)
mOPV	29 (11-43)	30 (0-66)
bOPV	41 (0-87)	45 (0-99)

Vaccine efficacy estimates % (95% CI)

Vaccine boycott in Nigeria

Special Report

OPV can wipe out a whole generation - Prof Kaita

Isa Sa'idu

WA: It is quite sometime since you raised the doubts that led to the controversy surrounding polio immunisation; have your doubts been cleared?
Professor Kaita: I am even surprised that you are here; I thought that this controversy has died down naturally because a lot has been said about it. I thought my contribution and other commentators were quite sufficient.

Professor Haruna Kaita is a staff in the faculty of Pharmaceutical Sciences, Ahmadu Bello University (ABU) Zaria. He has carried independent research on the polio vaccine and concluded that the vaccine is not safe.



Professor Haruna Kaita

...ve tested
 ...e found these
 ...ere you also able
 ...ily surface since
 ...ildren.
 ...that you should
 ...somebody no
 ...o ask when the

...lity in a seven
 ...angle. What
 ...is a risk to
 ...umaceutical
 ...matter how

...e as it is
 ...from 0-59
 ...ns if it is
 ...an entire
 ...that no

...ed and
 ...mple, it

there is a problem of generational development in the whole world especially in developing countries that are the victims.

In a nutshell, what we are saying is that these OPV are poisonous, those concerned should rectify them before anarchy surfaces especially in our country. We can't see danger and keep on going with it until everything is destroyed.

WT: Do you sense any conspiracy with the way the polio vaccination is being handled?

Kaita: I do not want to comment over this conspiracy theory because it is political while I am a scientist. I don't want to speculate. I want to pin these issues to hard facts and evidences as we did with the OPV. I have solid evidences with me that the OPV batch we analysed were contaminated.

NAFDAC has also tested these OPVs and

Special Report

Why we oppose polio vaccines - Nafi'u Baba-Ahmed

Nafi'u Baba Ahmed is the secretary general of the Supreme Council for Shari'a in Nigeria. In this interview with Weekly Trust, he gave reasons why his organisation oppose the oral polio immunisation exercise as being carried out.

Nasiru L. Abubakar

WT: What is your organisation's position on the ongoing polio immunisation?

Nafi'u Baba Ahmed: Our position has never changed. Our position remains as it is against the polio vaccine programme as it is today and as they are continuing tomorrow, because our basic fears have not been addressed.

As an organisation, the Supreme Council for Shari'ah in Nigeria, as a policy, has finished what it has to do on the issue of polio. We have undertaken the responsibility of drawing the attention of appropriate authorities- federal, state and local governments, and even individuals. We have sensitised people to be aware of the debilitating effect of the oral polio vaccines (OPV) as it is being done in Nigeria. We have finished.

All over the world today, the Supreme

is not so much an issue of concern. Our major concern has to do with the fact that, scientifically it has been tested and proven that there are some agents that are capable of preventing immunity. There are also some unknown contaminants. The federal government of Nigeria denied it that there were no contaminants. Then they accepted it that there are contaminants and there is also some level of these agents but that they are not in sufficient quantities to cause concern.

Still on the safety side, the World Health Organisation (WHO) that is at the fore-front, with UNICEF, World Bank and what have you, they have been saying that they have determined that a child has to be vaccinated, given OPV a number of times.

But in Nigeria some children are given 20 times more than that. Nobody has bothered to enquire as to the possible debilitating consequences in future. All they are concerned with is 'make sure everybody is given willingly with it'.

these diseases or in even helping parents to alleviate the cost of medical treatments, not even a kobo but they are expending billions upon billions on polio.

Look at, for instance (pointing to a newspaper report) EU spent \$12 billion- \$12 billion is about is more than one third of the total debt Nigeria was owing, for over 20 years we were worried about that debt. But EU alone is spending over \$12 billion on polio in Nigeria. But nobody will spend even \$1 million on measles or malaria. There is more to it than meets the eye.

Why polio? They have forced us to change the order of our priorities in terms of healthcare delivery. And why are they choosing polio vaccination because that can be applied to targeted countries. This is why in Nigeria, they are saying it is only in the north. Even in the north it is in the northwest. For example, they said there is one case in Kaduna, just one. But this is a matter that will be on CNN, BBC and so on when in the vein a thousand people may be killed by cerebrospinal meningitis and no body will show concern. These are food for thought for people to ponder.

Then there is this issue which is at the very root of our concern. In 1971, Henry Kissinger,



Nafi'u Baba Ahmed

available everywhere. Population control in whatever form is anti-Islamic. But it's not only Islam that is kicking against it. The Catholic Church the world over,

Nigeria - strategies for improving vaccine uptake



Immunisation Plus Days (IPDs)


- Community dialogue
- Improvements in coverage



Reduction in transmission in 2009


Pakistan & Afghanistan

- Shared borders areas and populations
- Security risks
- Days of tranquillity




President Karzai
Announces Polio Action
Group to Coordinate all
Ministries

Wild Poliovirus - 2011



Wild Poliovirus - 2012

01 January - 23 October



Excludes viruses detected from environmental surveillance and vaccine derived polioviruses. 1 WPV1 case in Gilgit Baltistan, date of onset 11 August 2012, does not appear on the map.

Explosive new outbreaks due to int'l spread



Tajikistan (Feb-Jul 2010)

458 cases

30% > 5 years of age




Congo (Oct 2010-Jan 2011)

476 cases

80% > 15 years of age

50% mortality

2^o spread - Central Asia & Caucasus 2010



Data in WHO HQ as of
15 Dec 2010

China 2011

21 cases between July-Oct 2011

11 cases are in adults 19-53 years of age



All cases from Xinjiang Uygur Autonomous Region of western China

Genetic sequencing → related to wild polioviruses circulating in Pakistan


Within 15 days:

- a 'level two' public health emergency had been declared
- both the Minister and Vice-Minister of Health had travelled to the affected region
- almost 150 health professionals from around China had been deployed
- five million doses of oral polio vaccine had been airlifted to the province
- more than 200,000 hospital records had been reviewed for potential polio cases

By October → 9 million children and young adults had been vaccinated

Eradication challenges - cVDPV (1)

- About 30% of immunized subjects excrete partially revertant strains of OPV called VDPV
- If immunization coverage is high, no problem
- However, at low levels of immunity VDPV can spread (i.e. $R_0 > 1$) and fully revert to wild type virulence (cVDPV)
- Leads to outbreaks of paralytic polio



From Kew et al. 2002


Madura Island, Indonesia - cVDPV (2005)




Eradication challenges - cVDPV (2)

- cVDPV outbreaks have now occurred in Hispaniola (2000-1), Philippines (2001), Madagascar (2001-2), China (2004), Indonesia (2005), Nigeria (2004-)...
- Retrospective studies have also identified cVDPV in Egypt (1988-93)
- Many cVDPV have been recombinants with wild enteroviruses
- Will be an important problem as countries scale-down immunization after polio considered to have been eliminated

Circulating Vaccine-derived Poliovirus*, Previous 6 Months**



Monthly distribution of cases by country



**10 Apr – 09 Oct 2012

Data in WHO HQ as of 09 October 2012


*Circulating Vaccine-derived poliovirus (cVDPV) is associated with 2 or more cases of AFP. VDPV type 2 cases with greater than or equal to 6nt difference from sabin in VP1; VDPV types 1 and 3 cases with greater than or equal to 10nt difference from sabin in VP1 are reported here. Figures exclude VDPV from non-AFP source. Figures may include different chains of transmission.

Additional eradication challenges


- Laboratory escape (esp. as present in stool sample etc. not collected for polio work)
- Factory escape (during manufacture of IPV)
- iVDPV - continued excretion of virus by vaccinated individuals
immuno-compromised individuals

Strategies for the cessation of OPV use (1)

- Abrupt (global OPV cessation day)
 - Excretion ceases after <6 months - environmental VDPV should disappear before enough new susceptibles
 - But, many possible sources for wildtype or VDPV introduction
 - Stockpile 850 million doses of monovalent OPV owned by WHO to control outbreaks by localised high coverage repeat immunisation
 - mOPV may itself reintroduce cVDPV depending on population movement



A new 'Endgame' strategy: parallel instead of sequential risk management



Strategies for the cessation of OPV use (2)

- Potential role for IPV
 - becoming vaccine of choice in wealthier nations (e.g. since 2004 used in UK), since risk of VAPP now outweighs that of wild poliovirus
 - Manufacture from Sabin poliovirus now being developed
- Problems
 - Must be injected
 - Low mucosal immunity (not yet demonstrated effective at preventing transmission in countries with poor sanitation)
 - Waning rather than life-long immunity in absence of circulating OPV?
 - Early vaccination schedule may have low seroconversion rates
 - Currently expensive, but enough to vaccinate all the world's children could be produced within 5-7 years

Further reading

- Grassly NC et al. New strategies for the elimination of polio from India. *Science* (2006) 314:1150-1153
- The Bulletin of the World Health Organization Special theme issue (2004) - Polio Eradication End-Stage Challenges Volume 82:1-81
- Minor PD Polio eradication, cessation of vaccination and re-emergence of disease *Nature Rev Microbiol* (2004) 2:473-482
- Aylward et al. in *The eradication of infectious diseases* (eds. Dowdle, W. R. & Hopkins, D. R.) p. 61-74 (John Wiley and Sons, Chichester, 1998).
- Kew O et al. Outbreak of poliomyelitis in Hispaniola associated with circulating type 1 vaccine-derived poliovirus. *Science* (2002) 296:356-359
- www.polioeradication.org