


# The Management of Diabetic Ketoacidosis in Adults



For young people under the age of 18 years use British Society of Paediatric Endocrinology and Diabetes (BSPED) guidelines: <http://www.bsped.org.uk/professional/guidelines/docs/DKAGuideline.pdf>

Diagnostic criteria: all three of the following must be present

- capillary blood glucose above 11 mmol/L
- capillary ketones above 3 mmol/L or urine ketones ++ or more
- venous pH less than 7.3 and/or bicarbonate less than 15 mmol/L

## BOX 1: Immediate management: time 0 to 60 minutes

(T=0 at time intravenous fluids are commenced)

If intravenous access cannot be obtained request critical care support immediately

**Action 1:** Commence 0.9% sodium chloride solution (use large bore cannula) via infusion pump.

See Box 2 for rate of fluid replacement

**Action 2:** Commence a fixed rate intravenous insulin infusion (IVI) (0.1 unit/kg/hr based on estimate of weight) 50 units human soluble insulin (Actrapid® or Humulin S®) made up to 50ml with 0.9% sodium chloride solution. If patient normally takes long acting insulin analogue (Lantus®, Levemir®) continue at usual dose and time

**Action 3:** Assess patient

- o Respiratory rate; temperature; blood pressure; pulse; oxygen saturation
- o Glasgow Coma Scale
- o Full clinical examination

**Action 4:** Further investigations

- Capillary and laboratory glucose
- Venous BG

- U & E
- FBC
- Blood cultures
- ECG
- CXR
- MSU

**Action 5:** Establish monitoring regimen

- Hourly capillary blood glucose
- Hourly capillary ketone measurement if available
- Venous bicarbonate and potassium at 60 minutes, 2 hours and 2 hourly thereafter
- 4 hourly plasma electrolytes
- Continuous cardiac monitoring if required
- Continuous pulse oximetry if required

**Action 6:** Consider and precipitating causes and treat appropriately

HDU/level 2 facility and/or insertion of central line may be required in following circumstances (request urgent senior review)

- Young people aged 18-25 years
- Elderly
- Pregnant
- Heart or kidney failure
- Other serious co-morbidities
- Severe DKA by following criteria
  - Blood ketones above 6 mmol/L
  - Venous bicarbonate below 5 mmol/L
- Venous pH below 7.1
- Hypokalaemia on admission (below 3.5 mmol/L)
- GCS less than 12
- Oxygen saturation below 92% on air (Arterial blood gases required)
- Systolic BP below 90 mmHg
- Pulse over 100 or below 60 bpm
- Anion gap above 16 [Anion Gap = (Na<sup>+</sup> + K<sup>+</sup>) - (Cl<sup>-</sup> + HCO<sub>3</sub><sup>-</sup>) ]

## BOX 2: Initial fluid replacement

Restoration of circulating volume is priority

**Systolic BP (SBP) below 90mmHg**

Likely to be due to low circulating volume, but consider other causes such as heart failure, sepsis, etc.

- Give 500ml of 0.9% sodium chloride solution over 10-15 minutes. If SBP remains below 90mmHg repeat whilst requesting senior input. Most patients require between 500 to 1000ml given rapidly.
- Consider involving the ITU/critical care team.
- Once SBP above 90mmHg give 1000ml 0.9% sodium chloride over next 60 minutes. Addition of potassium likely to be required in this second litre of fluid

**Systolic BP on admission 90 mmHg and over**

- Give 1000ml 0.9% sodium chloride over first 60 minutes

**Potassium replacement**

Potassium level (mmol/L)	Potassium replacement mmol/L of infusion solution
> 5.5	Nil
3.5-5.5	40 mmol/L
< 3.5	senior review – additional potassium required

## BOX 3: 60 minutes to 6 hours

**Aims of treatment:**

- Rate of fall of ketones of at least 0.5 mmol/L/hr OR bicarbonate rise 3 mmol/L/hr and blood glucose fall 3 mmol/L/hr
- Maintain serum potassium in normal range
- Avoid hypoglycaemia

**Action 1: Re-assess patient, monitor vital signs**

- Hourly blood glucose (lab blood glucose if meter reading 'HI')
- Hourly blood ketones if meter available
- Venous blood gas for pH, bicarbonate and potassium at 60 minutes, 2 hours and 2 hourly thereafter.
- If potassium is outside normal range, re-assess potassium replacement and check hourly. If abnormal after further hour seek immediate senior medical advice

**Action 2: Continue fluid replacement via infusion pump as follows:**

- 0.9% sodium chloride 1L with potassium chloride over next 2 hours
- 0.9% sodium chloride 1L with potassium chloride over next 2 hours
- 0.9% sodium chloride 1L with potassium chloride over next 4 hours
- Add 10% glucose 125ml/hr if blood glucose falls below 14 mmol/L

**More cautious fluid replacement in young people aged 18-25 years, elderly, pregnant, heart or renal failure. (Consider HDU and/or central line)**

**Action 3: Assess response to treatment**

Insulin infusion rate may need review if

- Capillary ketones not falling by at least 0.5 mmol/L/hr
- Venous bicarbonate not rising by at least 3 mmol/L/hr
- Plasma glucose not falling by at least 3 mmol/L/hr
- Continue fixed rate IVI until ketones less than 0.3 mmol/L, venous pH over 7.3 and/or venous bicarbonate over 18 mmol/L.

**If ketones and glucose are not falling as expected always check the insulin infusion pump is working and connected and that the correct insulin residual volume is present (to check for pump malfunction).**

If equipment working but response to treatment inadequate, increase insulin infusion rate by 1 unit/hr increments hourly until targets achieved.

**Additional measures**

- Regular observations and Early Warning Score (EWS)
- Accurate fluid balance chart, minimum urine output 0.5ml/kg/hr
- Consider urinary catheterisation if incontinent or anuric (not passed urine by 60 minutes)
- Nasogastric tube with airway protection if patient obtunded or persistently vomiting
- Measure arterial blood gases and repeat chest radiograph if oxygen saturation less than 92%
- Thromboprophylaxis with low molecular weight heparin
- Consider ECG monitoring if potassium abnormal or concerns about cardiac status

## BOX 5: 12 to 24 HOURS

**Expectation:** By 24 hours the ketonaemia and acidosis should have resolved. Request senior review if not improving

**Aim:**

- Ensure that clinical and biochemical parameters are continuing to improve or are normal
- Continue iv fluid replacement if not eating and drinking.
- If ketonaemia cleared and patient is not eating and drinking move to a variable rate IVI as per local guidelines
- Re-assess for complications of treatment e.g. fluid overload, cerebral oedema
- Continue to treat precipitating factors
- Transfer to subcutaneous insulin if patient is eating and drinking normally.

**Action 1 – Re-assess patient, monitor vital signs**

**Action 2 – Review biochemical and metabolic parameters**

- At 12 hours check venous pH, bicarbonate, potassium, capillary ketones and glucose
- Resolution is defined as ketones <0.3 mmol/L, venous pH>7.3
- If not resolved review fluid **Box 4 Action 1** and insulin infusion **Box 3 Action 3**

**Action 3**

If DKA resolved go to Box 6

## BOX 6: Resolution of DKA

**Expectation:** Patient should be eating and drinking and back on normal insulin.

If DKA not resolved identify and treat the reasons for failure to respond.

**This situation is unusual** and requires senior and specialist input.

**Transfer to subcutaneous insulin**

Convert to subcutaneous regime when biochemically stable (capillary ketones less than 0.3 mmol/L, pH over 7.3) and the patient is ready and able to eat. **Do not discontinue intravenous insulin infusion until 30 minutes after subcutaneous short acting insulin has been given**

Conversion to subcutaneous insulin should be managed by the Specialist Diabetes Team. If the team is not available use local guidelines. If the patient is newly diagnosed it is essential they are seen by a member of the specialist team prior to discharge.

Arrange follow up with specialist team.

## BOX 4: 6 to 12 hours

**Aims:**

- Ensure clinical and biochemical parameters improving
- Continue iv fluid replacement
- Avoid hypoglycaemia
- Assess for complications of treatment e.g. fluid overload, cerebral oedema
- Treat precipitating factors as necessary

**Action 1: Re-assess patient, monitor vital signs**

- If patient not improving by criteria in Box 3 seek senior advice
- Continue iv fluid via infusion pump at reduced rate
  - o 0.9% sodium chloride 1L with potassium chloride over 4 hours
  - o 0.9% sodium chloride 1L with potassium chloride over 6 hours
- Add 10% glucose 125ml/hr if blood glucose falls below 14 mmol/L

**Reassess cardiovascular status at 12 hours; further fluid may be required.**

**Check for fluid overload**

- **Action 2 – Review biochemical and metabolic parameters**
- At 6 hours check venous pH, bicarbonate, potassium, capillary ketones and glucose
- Resolution is defined as ketones less than 0.3 mmol/L, venous pH over 7.3 (do not use bicarbonate as a surrogate at this stage).
- Ensure referral has been made to diabetes team

**If DKA not resolved review insulin infusion (see BOX 3 Action 3)**

**If DKA resolved go to BOX 6**



**Groups represented:** Association of British Clinical Diabetologists; British Society for Endocrinology and Diabetes and Association of Children's Diabetes Clinicians; Diabetes Inpatient Specialist Nurse (DISN) Group; Diabetes UK; NHS Diabetes (England); Northern Irish Diabetologists; Society of Acute Medicine; Welsh Endocrine and Diabetes Society; Scottish Diabetes Group.